

# Pampas

*Cortaderia selloana*

## Family

Poaceae (grass)

## Also known as

Cutty grass, Prince-of-Wales feathers, common pampas

## Where is it originally from?

South America

## What does it look like?

Large-clump-forming grass (<4 m+). Leaf base is smooth or sparsely hairy, with no white waxy surface. Leaves are blueish-green above and dark green below, have a conspicuous midrib which does not continue into leaf base, no secondary veins between midrib and leaf edge, and snap readily when tugged. Dead leaf bases spiral like wood shavings. Erect, dense, fluffy, white-pinkish flowerheads (Jan-Jun) fade to a dirty white as seed forms.

## Are there any similar species?

*C. jubata* and native *Austroderia* species (toetoe) are similar. Toetoe leaves don't snap readily, midrib continues into leaf base, have distinct secondary veins between midrib and edge, and white waxy leaf sheaths. Dead leaves don't spiral. Drooping, light golden-yellow flowerheads (Sep-Jan).

## Why is it weedy?

Tolerates heat and frost, salt, wind, wet and drought, moderate-shade, most soils, low fertility, and recovers quickly after fire. Massive amounts of well dispersed seed are produced.

## How does it spread?

Seeds are spread very long distances by wind and occasionally water. Seeds are also spread by soil movement, dumped vegetation, contaminated forestry machinery, clothing, and on animal pelts. Common seed sources are plantation forests, roadsides, farm hedges, quarries, and wasteland.

## What damage does it do?

Colonises sprayed, burnt, slipped and otherwise disturbed sites and quickly becomes dense. Replaces groundcovers, shrubs, and ferns, creates fire hazards, provides habitats for possums and rats, and impedes access. Normally followed by weedy vines.

## Which habitats is it likely to invade?

Forest light gaps, slips, margins, disturbed sites, open habitats, riverbeds, cliffs, inshore and offshore islands, tussockland, fernland, herbfield, duneland, coastline, gumlands, salt marsh, estuaries, and shrublands.

## What can I do to get rid of it?

Establish that the species is not toetoe (look for erect seed heads in autumn. Toetoe begins flowering in summer and has drooping, golden coloured seed heads)

1. Physical removal - Hand pull, dig out or grub the plant to ensure all of the plant crown is removed.

2. Foliar spray early spring to mid-summer when plants are actively growing (and before the plants produce flowers in autumn)

Apply Glyphosate (360 g/L active ingredient) herbicide at 10ml/L plus organosilicone penetrant 2ml/L to cover entire plant. Note: Glyphosate overspray will kill other (desirable) plants.

OR apply Haloxypop herbicide (520g/l active ingredient at 3ml/L or 100g/l active ingredient at 15ml/L ) to cover all parts of plant but not to



[www.weedbusters.org.nz](http://www.weedbusters.org.nz)



Photo: Trevor James



Photo: Trevor James

the point it is dripping. Note: Haloxyfop herbicide is 'broadleaf friendly' but overspray will kill all grasses. Do not use over water bodies or wetlands and use only as directed on label.

*Tip* large areas of bare soil are ideal for this weed to establish. To avoid this, plant or sow desirable species to cover bare soil. Pampas seed can travel by wind over large distances. To contain spread, remove and dispose of flowerheads before they set seed.

**CAUTION:** When using any herbicide or pesticide, PLEASE READ THE LABEL THOROUGHLY to ensure that all instructions and directions for the purchase, use and storage of the product, are followed and adhered to.

### **What can I do to stop it coming back?**

Wind-born seeds will reinfest bare, burnt and sprayed sites. Grazed plants will resprout. Pampas is not a strong competitor at germination and prefers full sun, so encourage weed replacement (seeding, planting, regeneration) as you carry out control. Planting bare areas with broadleaf species will allow for a grass selective herbicide to be used. Dead pampas plants will burn easily. Plan for increased fire risk after control.



Photo: Trevor James